Rotation numbers in Thompson-Stein groups and applications

Isabelle LIOUSSE

arXiv:math/0604109v2 [math.DS] 31 Aug 2006

Abstract. We study the properties of rotation numbers for some groups of piecewise linear homeomorphisms of the circle. We use these properties to obtain results on PL rigidity, non isomorphicity, non exoticity of automorphisms, non smoothability for Thompson-Stein groups.

Keywords : PL homeomorphisms of the circle, rotation number, Thompson groups, automorphisms.
Mathematical subjet classification : 37E10, 37C15, 20E

1 Introduction.

In 1965, R. Thompson discovered the first example of a finitely presented infinite simple group. This group called T is defined as a group of PL homeomorphisms of the circle.

We recall that an homeomorphism f of the circle $S_r = \frac{\mathbb{R}}{\mathbb{Z}}$ of length r is **piecewise linear (PL)** if there exists a finite subdivision : $0 < a_1 < a_2 ...... < a_p = r$ such that
$\tilde{f}_{|[a_i,a_{i+1}]}(x) = \lambda_i x + \beta_i$, where $\tilde{f}$ is a lift of f to $\mathbb{R}$.

The a_i's are called **break points (or breaks)** of f, their set is denoted by $BP(f)$.

The λ_i's are called **slopes of** f**,** their set is denoted by $\Lambda(f)$.

The ratio $\sigma_f(a_i) = \frac{\lambda_i}{\lambda_{i-1}}$ is called the **jump** of f at a_i, their set is denoted by $\sigma(f)$.

The set of all orientation preserving (i.e. slopes are positive) PL homeomorphisms of S_r is denoted by $PL^+(S_r)$.

The **Thompson group** T consists of all orientation preserving homeomorphisms of $\frac{\mathbb{R}}{\mathbb{Z}}$ such that :
- breaks and their images are dyadic rational numbers,
- slopes are powers of 2.

The group T and its subgroup F –consisting of the elements of T that fix 0– occur in many areas of mathematics : logic, homotopy theory, group theory, dynamical systems. The standard survey for Thompson groups is [8]. Dynamical properties of T and F have been studied by Ghys and Sergiescu in [11] and Brin in [3].

In particular in [11], it is proved that :
- any non trivial representation $\phi : T \to Diff^r_+(S^1)$, $r \geq 2$ is semi-conjugate to the standard representation in $PL^+(S^1)$;
- T admits faithful representations in $Diff^\infty_+(S^1)$ and both of the two following situations can occur : all $\phi(T)$-orbits are dense or $\phi(T)$ has an exceptional minimal.

As corollary and using Denjoy theorem, it's shown in [11] that the rotation number of any element of T is rational.

In [12], É. Ghys asked about the following problem : consider a piecewise linear homeomorphism f of S^1 such that all its slopes, breaks, images of breaks are rational numbers, is it true that the rotation number of f is rational ?
This question has a negative answer. Counterexamples can be found in [2], a Boshernitzan paper and not explicitly stated by Herman in [13] chapt VI section7.

In [3], M. Brin proved that the outer automorphism group of T has order 2, as is the group $OUT(Homeo^+(S^1))$. This fact confirms the fact that the finitely generated group T is imitated the behaviors of the continuous group $Homeo^+(S^1)$ (simplicity, cohomology, automorphisms).

An other interesting property of T and F due to Farley is that T and F are a-T-menable (see [10]). It's still unknown if F is amenable or not.

Moreover, generalisations of Thompson groups have been quite a lot defined and studied : Higman [14], Bieri-Strebel [1], Brown [7], Stein [22] and Brin-Guzman ([6]). In this paper, we consider the generalisations in $PL(S^1)$ and follow the descriptions of Bieri-Strebel and Stein.

Thompson-Stein groups.

Definitions. Let $\Lambda \subset \mathbb{R}^{+*}$ be a multiplicative subgroup, and $A \subset \mathbb{R}$ be an additive subgroup, invariant by multiplication by elements of Λ.
We define $T_{r,\Lambda,A}$ as the subgroup of $PL^+(S_r)$ consisting of elements with slopes in Λ, breaks and their images in A. The slope set of $T_{r,\Lambda,A}$ is Λ and the break set of $T_{r,\Lambda,A}$ is A.

In the case $\Lambda =< n_i >$ the multiplicative group generated by p integers $1 < n_1 < ... < n_p$ independent (in the sense of free abelian group generators, this is equivalent to the statement that the $\log n_i$ are $\mathbb{Q}$-independent) and $A = \mathbb{Z}[\frac{1}{n_1}, ..., \frac{1}{n_p}] = \mathbb{Z}[\frac{1}{m}]$ where $m = lcm(n_i)$ the least common multiple of the n_i :

the group $T_{r,\Lambda,A}$ is denoted by $T_{r,(n_i)}$ and called a **Thompson-Stein group**. It's the smallest non trivial subgroup $T_{r,<n_i>,A}$. The subgroup of $T_{r,(n_i)}$ consisting of elements that fix 0 is denoted by $F_{r,(n_i)}$.

When $p = 1$, the group $T_{r,(m)}$ [resp. $F_{r,(m)}$] is denoted by $T_{r,m}$ [resp. $F_{r,m}$]. The classical Thompson groups are $T = T_{1,2}$ and $F = F_{1,2}$.

Questions. For these groups many natural questions arise :
How are the rotation numbers of the elements of $T_{r,(n_i)}$?
How are automorphisms of $T_{r,(n_i)}$?
Is it possible to describe all $T_{r,(n_i)}$ up to isomorphisms, up to quasi isometry ?
Is it possible to classify the representations of $T_{r,(n_i)}$ in $PL^+(S_r)$, $Diff^k_+(S_r)$? In particular, are the $T_{r,(n_i)}$ (infinitely) smoothable ?

Properties. M. Stein in [22] proved that :
- $T_{r,(n_i)}$ and $F_{r,(n_i)}$ are finitely presented and FP^∞,
- $T''_{r,(n_i)}$ and $F'_{r,(n_i)}$ are simple groups,
- precise homology informations.

In [1] –the author is attached to mention that she couldn't get this unpublished paper, so she refered to results that are cited or reproved in [22], [3] and [6]– Bieiri and Strebel have studied isomorphisms and automorphisms of $T_{r,\Lambda,A}$.

They've described PL isomorphisms between distincts $T_{r,(n_i)}$; in particular, they proved that : " $T_{r,(n_i)}$ and $T_{r',(n_i)}$ are PL-isomorphic if and only if , $(r - r')$ is an integer multiple of $d = gcd(n_i - 1)$ the highest common divisor of the $n_i - 1$".

They also proved that automorphisms of $F_{r,\Lambda,A}$ are not exotic (i.e. they are realized as PL conjugation) provided Λ is a dense multiplivative group.

To extend this result to $T_{r,(n_i)}$ (in particular for $p \geq 2$), it's reasonnable to think about localization results (such results can be found in [3] and [6] for $T_{r,m}$ when r is not a multiple of $d = m - 1$). A very surprising theorem is proved in [6] : $T_{m-1,m}$ has (infintely many) exotic automorphisms.

Here, we developp a different approach to study the automorphisms of Thompson-Stein groups : we analyse the rotation numbers and use a rigidity result for actions in $PL^+(S^1)$ or $Diff^k_+(S^1)$ to obtain as corollaries informations on isomorphisms and automorphisms.

Theorem 1. Rational rotation numbers. *Let $m \geq 2$, $r \geq 1$ and $q \geq 1$ be integers. Then*

A. Each element of $T_{r,<m>,\mathbb{Q}}$ has rational rotation number, so has periodic points.

B. The following properties are equivalent :
(1) there exists $p \in \mathbb{N}^*$ such that $T_{r,m}$ contains an element of reduced rotation number $\frac{p}{q}$,
(1') for all $p \in \mathbb{N}^*$, the group $T_{r,m}$ contains an element of reduced rotation number $\frac{p}{q}$,
(2) gcd $(m-1, q)$ is a divisor of r,
(3) the group $T_{r,m}$ contains an element of order q
(3') for all $p \in \mathbb{N}^*$, the group $T_{r,m}$ contains an element of order q with rotation number $\frac{p}{q}$.

Immediate consequences of Theorem 1. Fix two integers $m \geq 2$ and $r \geq 1$.
On the rotation number map $\rho :$ Homeo$^+(S_r) \to \frac{\mathbb{R}}{\mathbb{Z}}$ defined by $\rho(f) = \lim_{n\to\infty} (\tilde{f}^n(0)/rn)\ (mod 1)$.

1. The image set $\rho(T_{r,m}) \subset \frac{\mathbb{Q}}{\mathbb{Z}}$ and any number in $\rho(T_{r,m})$ is realized as the rotation number of a finite order element of $T_{r,m}$.

2. Any rational number is realized as the rotation number of an element of finite order of $T_{m-1,m}$; in particular, $\rho(T_{m-1,m}) = \frac{\mathbb{Q}}{\mathbb{Z}}$.

3. The group $T_{m-1,m}$ is the only group for $0 < r < m-1$ that contains elements of order $m-1$. In particular, $\rho(T_{r,m}) \neq \frac{\mathbb{Q}}{\mathbb{Z}}$ for $0 < r < m-1$ (mod $m-1$).

4. If m, r are odd, the group $T_{r,m}$ doesn't contain elements of order 2 (of any even order).

5. The image set $\rho(T_{1,m}) = \{\frac{p}{q} : \gcd(q, m-1) = 1\}$. If $m-1$ is a prime number then $\rho(T_{1,m}) = \rho(T_{r,m})$, for all $0 < r < m-1$ (mod $m-1$).

On the isomorphic problem.

6. The group $T_{m-1,m}$ is isomorphic to none of the groups $T_{r,m}$ with $0 \leq r < m-1$.

7. If m is odd, r odd and r' even, the groups $T_{r,m}$ and $T_{r',m}$ are not isomorphic.

In the next theorem, we prove that Thompson-Stein groups contain free abelian groups that act freely on S_r. The rank of such a group depends on the rank of the slope set.

Theorem 2. Irrational rotation numbers. *Let $G = T_{r,(n_i)}$ be a Thompson-Stein group associated to $p \geq 1$ independent integers n_i. Then :*

A. The group G contains a free abelian group of rank $p-1$ acting freely. That is equivalent to the statement that G contains $(p-1)$ commuting homeomorphisms with irrational rotation numbers ρ_i such that 1 and the ρ_i are $\mathbb{Q}$-independent.

B. The group G doesn't contain any free abelian group of rank p acting freely on the circle.

C. The rotation number of any element in a free abelian subgroup of rank 2 of G is written in the form $\rho(g) = \frac{\log\alpha}{\log\beta}$, with $\alpha, \beta \in < n_i >$.

C'. Any number $\frac{\log\alpha}{\log\beta} \in [0, 1[$ with $\alpha, \beta \in < n_i >$ can be realized as the rotation number of an element of $T_{d,(n_i)}$, where $d = gcd(n_i - 1)$.

Consequences of theorem 2. *Let $G = T_{r,(n_i)}$ and $A = \mathbb{Z}[\frac{1}{m}]$, with $m = lcm(n_i)$.*

1. The free rank of G defined by $frk(G) = \max\{n \in \mathbb{N} : \exists \mathbb{Z}^n \subset G$ acting freely$\} = p-1$.

2. Any representation ϕ from $G = T_{r,(n_i)}$, $p \geq 3$ into $PL^+(S^1)$ that is topologically conjugate to the standard representation in $PL^+(S_r)$ is PL-conjugate to the standard representation.

3. The group $G = T_{r,(n_i)}$, $p \geq 3$ has no exotic automorphism ; more precisly, its automorphisms are in realized by conjugation by maps in $T_{r,invA,A}$, where $invA = \{a \in A : \frac{1}{a} \in A\}$. In particular, OUTG has order 2 provided the n_i are coprime.

4. Let $r \in \mathbb{N}^*$, two distincts Thompson-Stein groups $T_{r,(n_1,\dots n_p)}$ with $p \geq 3$ and $T_{r,(n'_1,\dots,n'_q)}$ with $q \geq 1$ are not isomorphic.

Remarks on theorem 2 and aknowlegments. According to G. Rhin ([21]), the $\frac{\log\alpha}{\log\beta}$ are diophantine numbers. I'm grateful to Yann Bugeaud and Nicolas Brisebarre for indicating me references and details on this property.

The proof of theorem 2A is based on extensions of Boshernitzan examples. Part B and C are consequences of a rigidity result for free PL-actions of $\mathbb{Z}^2$ on the circle due to Minakawa ([18]) and the explicit calculation of rotation numbers for some particular PL-homeomorphisms.

Theorem 3 (Representations in $PL(S^1)$ or $Diff(S^1)$).

Let $G = T_{1,(n_i)}$ be the Thompson-Stein group associated to $p \geq 2$ independent integers : $(2, n_2, ..., n_p)$. Then :

A. Each non trivial [faithfull] representation ϕ from G into $PL(S^1)$ or $Diff^2(S^1)$ is topologically conjugate to the standard representation in $PL(S_r)$.

B. There exists $k \geq 2$ depending on the $logn_i$'s diophantine coefficients such that any representation from G into $Diff^k(S^1)$ is trivial [has finite image]. In particular, G is not realizable in $Diff^\infty(S^1)$.

The proof of theorem 3 is the combination of :
- dynamics properties of G and its subgroups acting on S^1, following Ghys-Sergiescu approach. It's important to say that the theorem K in [11] holds for PL-actions instead of $Diff^2(S^1)$ (as Koppel lemma holds for PL-actions) and doesn't extend to $T_{r,m}$ with $m > 2$ since the Euler class triviality argument doesn't work with $rk(H^2(T_{r,m}, \mathbb{Z}) > 2$.
- and the Brin version ([4]) of a Rubin's theorem ([21]).

Consequence of theorems 2 and 3. *Let $G = T_{1,(n_i)}$ be a Thompson-Stein group associated to $p \geq 3$ independant integers : $(2, n_2, ..., n_p)$. Any non trivial representation ϕ from G into $PL(S^1)$ is PL-conjugate to the standard representation.*

A important piece for our proofs is the Bieri-Strebel criterion.

Definition. Two real intervals $[a, c]$ and $[a', c']$ are $PL_{\Lambda,A}$-**equivalent** if there exists a PL homeomorphism with slopes in Λ and breaks in A that takes $[a, c]$ to $[a', c']$.

Bieri-Strebel criterion. The intervals $[a, c]$ and $[a', c']$ are $PL_{\Lambda,A}$-equivalent if and only if their length are equal modulo $(1-\Lambda)A$, where the set $(1-\Lambda)A = \{\sum(1-\lambda_i)a_i, \lambda_i \in \Lambda, a_i \in A\}$ (for a proof, see [22] appendix).

For Thompson-Stein groups. Suppose that $\Lambda =< n_i >$ and $A = \mathbb{Z}[\frac{1}{m}]$ with $m = lcm(n_i)$, then :
- the set $(1 - \Lambda)A = dA = (d\mathbb{Z})\Lambda$, where $d = gcd(n_i - 1)$.
- two integers are equal modulo $(1 - \Lambda)A$ iff their difference is a multiple of d.
- the groups $T_{r,(n_i)}$ and $T_{r',(n_i)}$ are PL-conjugate provided $r = r'$ (mod d).

Proof of $(1 - \Lambda)A = dA$.

Step 1. $(1-\Lambda)A \subset dA$. As $(1-\Lambda)A$ is a submodule, it suffices to prove that $(1-\lambda)a \in dA$, with $a \in A$ and $\lambda = n_1^{s_1}...n_p^{s_p} \in \Lambda$. By reducing to same denominator, $(1 - \lambda)a = (1 - n_1^{a_1}...n_p^{a_p})a = (n_1^{b_1}...n_p^{b_p} - n_1^{c_1}...n_p^{c_p})a'$ with b_i, c_i positive integers, $a' \in A$. By replacing the n_i by $k_id + 1$ and developping, we get $(1 - \lambda)a = (dN)a'$ with $N \in \mathbb{N}$.

Step 2. $dA \subset (1 - \Lambda)A$. Let $a \in A$, using Bezout identity we get $d = u_1(n_1 - 1) + ... + u_p(n_p - 1)$ and hence $da = \sum(n_i - 1)(u_ia) \in (1 - \Lambda)A$.

2 Proof of Theorem 1.

Proof of theorem 1.A. The first step is to prove that each $f \in T_{r,m}$ has rational rotation number. The homeomorphism $f \in T_{r,m}$ of $S_r = \frac{[0,r]}{0=r}$ is identified with the bijection $\tilde{f}$ (mod r) of $[0,r[$, where $\tilde{f}$ is a lift to $\mathbb{R}$ of f.

For all $x \in [0,r[$, we have $f(x) = m^{e(x)}x + \dfrac{p(x)}{m^{k(x)}}$, with $e(x) \in \mathbb{Z}$, $p(x) \in \mathbb{Z}$ and $k(x) \in \mathbb{N}$. Note that these three functions are bounded by constants independent of x.

The orbit of 0 contains only m-adic numbers, that is for all $n \in \mathbb{Z}$, the point $f^n(0)$ is written is the form $\quad f^n(0) = \dfrac{M_n}{m^{N_n}}$, $\quad$ with $N_n \in \mathbb{N}$ and $(*)$ $\; M_n \in \mathbb{N}$ is not a multiple of m or is zero. Remark that if m is not a prime number, the fraction may be unreduced, but the condition $(*)$ assumes the uniqueness of the expression $f^n(0) = \frac{M_n}{m^{N_n}}$.

If there exists a non zero integer n such that $M_n = 0$ then $f^n(0) = 0$, therefore the orbit of 0 is periodic and the rotation number of f is rational, we are done. In what follows, we assume that for all positive integer n the integer M_n is not a multiple of m.

Fact 1 : *$N_n \to +\infty$ when $n \to +\infty$ or the orbit of 0 is periodic.*

By absurd, if the sequence N_n doesn't tend to $+\infty$, there exists a bounded subsequence N_{s_n}, we write $N_{s_n} \leq B$. In this case, the subsequence $f^{s_n}(0)$ is contained in a finite set : $\{\dfrac{p}{m^k} \;$ with $\; 0 \leq k \leq B \;$ and $\; 0 \leq p \leq r.m^B\}$. But, this is possible iff the orbit of 0 is periodic.

Fact 2 : *If $N_n \to +\infty$ then $Df^n(0) \to 0$ when $n \to +\infty$.*

We suppose that $N_n \to +\infty$ and compute $f^{n+1}(0)$:

$f^{n+1}(0) = f(f^n(0)) = m^{e(f^n(0))}\dfrac{M_n}{m^{N_n}} + \dfrac{p(f^n(0))}{m^{k(f^n(0))}} = \dfrac{M_n}{m^{N_n - e(f^n(0))}} + \dfrac{p(f^n(0))}{m^{k(f^n(0))}}$.

As N_n tends to $+\infty$, as e and k are bounded, there is an integer n_0 such that for all $n \geq n_0$, we have $N_n - e(f^n(0)) > k(f^n(0))$. It follows that :

$$f^{n+1}(0) = \frac{M_n + m^{N_n - e(f^n(0)) - k(f^n(0))}p(f^n(0))}{m^{N_n - e(f^n(0))}} = \frac{M_{n+1}}{m^{N_{n+1}}}.$$

As m is a divisor of $m^{N_n - e(f^n(0)) - k(f^n(0))}$ but not a divisor of M_n, the integer m is not a divisor of $M_n + m^{N_n - e(f^n(0)) - k(f^n(0))}p(f^n(0))$. The uniqueness of the expression $f^{n+1}(0) = \frac{M_{n+1}}{m^{N_{n+1}}}$, implies that $N_{n+1} = N_n - e(f^n(0))$, for all $n \geq n_0$.

Thus, for all $p \in \mathbb{N}^*$, we have $N_{n_0+p} = N_{n_0} - \left(e(f^{n_0}(0)) + ... + e(f^{n_0+p-1}(0))\right)$. Hence, $e(f^{n_0}(0)) + ... + e(f^{n_0+p-1}(0)) \to -\infty$ when $p \to +\infty$. Finally, $e(0) + e(f(0)) + ... + e(f^n(0)) \to -\infty$ when $n \to +\infty$. But $Df^n(0) = m^{e(0) + e(f(0)) + ... + e(f^n(0))}$ so $Df^n(0) \longrightarrow 0$.

Conclusion. If the rotation number of f is irrational :

- The f-orbit of 0 is not periodic. From this and fact 1 follows that $N_n \to +\infty$ and thus using fact 2 we get that $Df^n(0) \to 0$.

- Denjoy's inequalities apply to f and give $Df^{q_n}(0) \geq e^{-Var log Df}$ that can't tend to 0.

From this contradiction follows the rationality of the rotation number of $f \in T_{r,m}$.

End of the proof of theorem 1.A. The group $T_{r,<m>,\mathbb{Q}}$ is identified with a group of piecewise linear bijections of $[0,r[$ with slopes power of m, breaks and images of breaks rational numbers. Let $f \in T_{r,<m>,\mathbb{Q}}$, on each continuity interval of Df, we write $f(x) = m^{e(x)}.x + \frac{p(x)}{q(x)}$. Let Q be the lcm of the $q(x)$, the homothety $\mathcal{H}_Q : S_r \to S_{Qr}$ $(\mathcal{H}_Q(x) = Qx)$ conjugate f to an element of $T_{Qr,m}$ for which the first part applies.

Proof of theorem 1.B.

Easy implications.

(3) $\Rightarrow$ (1) is clear, since f of order q has rotation number $\rho(f) = \frac{p}{q}$ with $(p, q) = 1$.

(1) $\Leftrightarrow$ (1′) and (3) $\Leftrightarrow$ (3′) are obtained by choosing suitable powers of f.

We now prove the two implications (1) $\Rightarrow$ (2) $\Rightarrow$ (3)**.**

(1) $\Rightarrow$ (2). By hypothesis, there exists $f \in T_{r,m}$ with $\rho(f) = \frac{p}{q}$. According to Poincaré, f has a periodic point a of period q. Replacing f by a suitable power, we may assume $\rho(f) = \frac{1}{q}$.

Fix $\tilde{f}$ a lift of f to $\mathbb{R}$ and identify f with the bijection $\tilde{f}(mod\ r)$ of $[0, r[$. By replacing a by its nearest from 0 iterate, the orbit of a can be ordered as :

$$0 \leq a = a_0 < f(a) = a_1 < .... < f^{q-1}(a) = a_{q-1} < r$$

The intervals $[0, a]$ and $[f(0), f(a)]$ are $PL_{\Lambda,A}$-equivalent so, using the Bieri-Strebel criterion, $a = (f(a) - f(0))\ mod(1 - \Lambda)A$ and thus $f(a) - a = f(0)\ mod(1 - \Lambda)A$.

Furthermore, (via the map $\tilde{f}$) the intervals $[a_0, a_1]$, $[a_1, a_2]$, ... and $[a_{q-1}, a_0 + r]$ are $PL_{\Lambda,A}$-equivalent so their lengths l_i all equal $mod(1 - \Lambda)A$ the length of the interval $[a_0, a_1]$ that is $f(a) - a$.

Finally, the lengths l_1, l_2,....l_q of the intervals $[a_0, a_1]$, $[a_1, a_2]$, ...,$[a_{q-1}, a_0 + r]$ equal $f(0)$ $mod\ (1 - \Lambda)A = dA$. Adding $l_1 + l_2 + ... + l_q$, we obtain $r = qf(0)\ mod\ dA$, so $r - qf(0) \in (m-1)\mathbb{Z}[\frac{1}{m}]$. Thus, there exists integers u, v, s such that $r - qf(0) = (m-1)\frac{v}{m^s}$ and $f(0) = \frac{u}{m^s}$, so such that $m^s r - qu = (m - 1)v$, that is to say $m^s r = qu + (m - 1)v$. This implies that $m^s r$ is a multiple of $gcd(q, m - 1)$. But $(m - 1)$ and m (also m^s) are coprime, we conclude therefore that r is a multiple of $gcd(q, m - 1)$.

(2) $\Rightarrow$ (3). Fix r, m two positive integers and suppose that r is a multiple of $gcd(q, m - 1)$. Using Bezout, $r = uq + v(m - 1)$, so $r = uq$ modulo $(m - 1)$. The Bieri-Strebel criterion implies that the groups $T_{uq,m}$ and $T_{r,m}$ are isomorphic. Furthermore, the group $T_{uq,m}$ contains the rotation $x \mapsto x + u$ of order q. This completes the proof.

3 Proof of theorem 2 and consequences.

The proof of theorem 2.A is based on generalisations of the examples gived by Boshernitzan in [2]. The first type of generalisation is to pass from the usual circle to S_r and the second is to construct carefully examples in order to obtain break points in the suitable set A and commuting examples.

3.1 Boshernitzan examples on S_r.

Consider $S_r = \frac{\mathbb{R}}{r\mathbb{Z}} = \frac{[0,r]}{0=r}$ the circle of length $r \in \mathbb{N}^*$ and $f \in PL^+(S_r)$ with exactly two break points a and $f(a) = 0$. The map f is identified with the bijection $\tilde{f}$ (mod r) of $[0, r[$.



Fig. 1

Fixing the initial break to be 0, such a map is uniquely defined by its slopes λ_1 and λ_2. Then, it admits 0 , $a = r\frac{1-\lambda_2}{\lambda_1-\lambda_2}$ as breaks and 0 , $b = f(0) = 1 - \lambda_1 a$ as images of breaks.

Conversely, given two positive numbers λ_1 and λ_2 such that $(\log\lambda_1 \log\lambda_1) < 0$, there exists a unique $f \in PL^+(S_r)$ with breaks 0 and a and slopes λ_1 and λ_2 such that $f(a) = 0$. This PL homeomorphism f is denoted by $f_{r,\lambda_1,\lambda_2}$. Let's define :

- the homothety $\mathcal{H}_r : \begin{cases} S^1 = S_1 & \to S_r \\ x & \mapsto rx \end{cases}$ of ratio r and
- h_σ the homeomorphism of S^1 identified with the restriction to $[0,1[$ of one of its lift :

$$h_\sigma(x) = \frac{\sigma^x - 1}{\sigma - 1} \quad \text{if } \sigma \in]0,1[\qquad \text{and} \qquad h_1(x) = x.$$

It's easy to check (by an explicit calculation, as it's done in [2]), that :

▷ the map $\mathcal{H}_r \circ h_{\frac{\lambda_1}{\lambda_2}}$ conjugates $f_{r,\lambda_1,\lambda_2}$ to the rotation of S^1 of angle $\frac{\log\lambda_1}{\log\lambda_1 - \log\lambda_2}$,

▷ in particular, f has rotation number $\rho(f) = \dfrac{\log\lambda_1}{\log\lambda_1 - \log\lambda_2} (mod\ 1)$.

▷ Conversely, for all $\rho \in S^1$, for all $\sigma \in]0,1[$, the map $(\mathcal{H}_r \circ h_\sigma) \circ R_\rho \circ (\mathcal{H}_r \circ h_\sigma)^{-1}$ is in $PL^+(S_r)$ and has two breaks 0 and a satisfying $f(a) = 0$, it's called a **Boshernitzan on S_r.**

The map defined by $B_\rho^\sigma = h_\sigma \circ R_\rho \circ h_\sigma^{-1}$ for $\sigma \in]0,1]$ is a classical Boshernitzan on S^1.

To sum up, $f_{r,\lambda_1,\lambda_2} = (\mathcal{H}_r \circ h_\sigma) \circ R_\rho \circ (\mathcal{H}_r \circ h_\sigma)^{-1} = \mathcal{H}_r \circ B_\rho^\sigma \circ \mathcal{H}_r^{-1}$,

where (λ_1, λ_2) and (ρ, σ) are related by $\begin{cases} \sigma = \frac{\lambda_1}{\lambda_2} \\ \rho = \frac{\log\lambda_1}{\log\lambda_1 - \log\lambda_2} \end{cases}$ and $\begin{cases} \lambda_1 = \sigma^{\rho-1} \\ \lambda_1 = \sigma^\rho \end{cases}$

It's easy to check that two Boshernitzan of irrational rotation numbers commute if and only if they have the same jump $\sigma = \frac{\lambda_1}{\lambda_2}$ at 0, since they are conjugate to rotations through the same normalized homeomorphism.

3.2 The Ghys question and remarks.

Let's n_1 and n_2 be two independents integers. Consider f the Boshernitzan of S^1 with slopes $\lambda_1 = n_1$ and $\lambda_2 = \frac{1}{n_2}$. The data of f are rational. Thus $f \in T_{1,\mathbb{Q},\mathbb{Q}}$ but its rotation number $\rho(f) = \frac{\log n_1}{\log n_1 + \log n_2} \notin \mathbb{Q}$. In general, because of the denominator in the formula that gives the break point $a = \frac{1-\lambda_2}{\lambda_1-\lambda_2}$, the previous PL homeomorphism is not in the Thompson-Stein group $T_{1,(n_1,n_2)}$. For example, by taking $n_1 = 2$, $n_2 = 3$, the associated break point is $a = \frac{2}{5} \notin \mathbb{Z}[\frac{1}{6}]$. Note that, for proving that a Boshernitzan f of S_r is in $T_{r,\Lambda,A}$, it suffices to prove that its slopes λ_1 and λ_2 are in Λ and that the break a is in A (from this follows that $f(0) \in A$ because $f(0) = r - \lambda_1 a = \lambda_2(r - a)$). The examples with irrational rotation numbers that we will construct in Thompson-Stein groups are not Boshernitzan examples they are PL conjugate to Boshernitzan examples.

3.3 Proof of theorem 2.A.

Here, we construct elements of irrational rotation numbers in any Thompson-Stein group having slope set of rank p at least 2. Let $G = T_{r,(n_1,\ldots,n_p)}$ be a Thompson-Stein group, $m = lcm(n_i)$ and $d = gcd(n_i - 1)$.

Step 1 : there exists $\Pi \in \Lambda$, $\Pi = n_1^{\alpha_1} n_2^{\alpha_2} \ldots n_p^{\alpha_p}$ with $\alpha_i \in \mathbb{N}^*$ such that $gcd(\frac{\Pi-1}{d}, d) = 1$.

By definition of d, we have $n_i = k_i d + 1$ where k_i are p positive coprime integers. So there exists p integers $(\beta_1, \ldots., \beta_p)$ such that $\quad k_1\beta_1 + \ldots. + k_p\beta_p = 1 \quad (*)$.

Fix p positive integers $\alpha'_i \in \mathbb{N}^*$ such that $\alpha'_i > |\beta_i|$.

If $k_1\alpha'_1 + ... + k_p\alpha'_p$ and d are coprime then we set $\alpha_i = \alpha'_i$.

If not, $k_1\alpha'_1 + ... + k_p\alpha'_p = w'd'$ with $d' > 0$ a divisor of d. By multiplicating this equality by $n = \frac{d}{d'}$, we obtain : $k_1(n\alpha'_1) + ... + k_p(n\alpha'_p) = w'd$. Adding (*) to this equality, we get $k_1(n\alpha'_1 + \beta_1) + ... + k_p(n\alpha'_p + \beta_p) = w'd + 1$ and we set $\alpha_i = n\alpha'_i + \beta_i > 0$. It's clear that d is relatively prime to $w'd + 1$.

Hence in both cases, we have found p positive integers $\alpha_i > 0$ such that $w = k_1\alpha_1 + ... + k_p\alpha_p$ and d are coprime. Consider $\Pi = n_1^{\alpha_1} n_2^{\alpha_2} ... n_p^{\alpha_p}$ and compute $\Pi - 1$. We denote by C_n^p the binomial coefficient $C_n^p = \frac{n!}{p!(n-p)!}$.

$$\Pi - 1 = (k_1 d + 1)^{\alpha_1} ....(k_p d + 1)^{\alpha_p} = \sum_{\substack{i_j \in \{0, ..., \alpha_j\} \\ j \in \{1, ...p\}}} (C_{\alpha_1}^{i_1} ... C_{\alpha_p}^{i_p})(k_1^{i_1} .... k_p^{i_p}) d^{(i_1 + ... i_p)} \quad -1 =$$

$$\sum_{\substack{i_j \in \{0, ..., \alpha_j\}, \sum i_j \geq 2 \\ j \in \{1, ...p\}}} (C_{\alpha_1}^{i_1} ... C_{\alpha_p}^{i_p})(k_1^{i_1} .... k_p^{i_p}) d^{(i_1 + ... i_p)} \quad + \sum_{\substack{i_j \in \{0, ..., \alpha_j\}, \sum i_j \leq 1 \\ j \in \{1, ...p\}}} \quad ..... \quad -1 =$$

$$= d^2\Sigma + d(k_1\alpha_1 + ... + k_p\alpha_p) = d^2\Sigma + dw \;\text{ where }\; \Sigma \in \mathbb{N}.$$

Because, $\sum i_j \leq 1$ if and only if all i_j are zero (in this case the term of the sum is 1) , or all i_j are zero except one i_l that is egal to 1 (in this case the term of the sum is $\alpha_l k_l d$).

Using that $w = \frac{\Pi - 1}{d} - d\Sigma$ and d are coprime, we conclude that $\frac{\Pi - 1}{d}$ and d are coprime.

Step 2 : Construction of maps with irrational rotation numbers.

Fix $j \in \{1, ..., p\}$ and set $r_k = k\frac{\Pi - 1}{d}$ for $k \in \mathbb{N}^*$, with Π given by the previous step.
Fix $i \in \{1, ..., p\}$ and consider f_i the Boshernitzan on S_{r_k} defined by its two slopes :

$$\lambda_1 = n_i > 1 \text{ and } \lambda_2 = \frac{\lambda_1}{\Pi} < 1.$$

For proving that f is in $T_{r_k,(n_i)}$, it suffices to prove that its break $a_i \in \mathbb{Z}[\frac{1}{m}]$. We have $a_i = r_k \dfrac{1 - \lambda_1}{\lambda_2 - \lambda_1} = k(\dfrac{\Pi - 1}{d}) \times \dfrac{1 - \lambda_1}{\lambda_1 \Pi^{-1} - \lambda_1} = \dfrac{k\Pi}{\lambda_1} \dfrac{\lambda_1 - 1}{d}$. But, $1 - \lambda_1 \in dA$ so $\frac{\lambda_1 - 1}{d} \in A$, thus $a_i \in A$.

To sum up : for any $i \in \{1, ..., p\}$, we have construct $f_i \in T_{r_k,(n_i)}$ such that :

$$\rho(f_i) = \frac{\log n_i}{\log \Pi} \notin \mathbb{Q} \quad \text{and} \quad \sigma_{f_i}(0) = \Pi.$$

Futhermore, for $i \in \{2, ..., p\}$, the f_i have irrational rotation numbers such that $\rho(f_i)$ and 1 are $\mathbb{Q}$-independent, and commute because have the same jump $\sigma_{f_i}(0) = \Pi$. Therefore the Thompson-Stein group $T_{r_k,(n_i)}$ contains a $\mathbb{Z}^{p-1}$ that acts freely on S^1.

The classes in $\frac{\mathbb{Z}}{d\mathbb{Z}}$ described by the $r_k = k\frac{\Pi - 1}{d}$, when k ranges over $\mathbb{N}^*$ are the multiples of the class in $\frac{\mathbb{Z}}{d\mathbb{Z}}$ of $gcd(d, \frac{\Pi - 1}{d}) = 1$. So any positive integer r is in the class ($mod\ d$) of some r_k. By Bieri-Strebel criterion, the group $T_{r,(n_i)}$ is PL-conjugate to some $T_{r_k,(n_i)}$, thus it contains a $\mathbb{Z}^{p-1}$ that acts freely on S^1. This ends the proof of theorem 2, part A.

3.4 Proof of theorem 2 B, C and C'.

These proofs use a rigidity result for PL-actions of $\mathbb{Z}^2$ on circles, which is a lemma in [18] : *"If $f, g \in PL^+(S_r)$ generate a freely acting group then f and g are PL conjugate to two commuting Boshernitzan on S_r"*. Here, we slightly modified the proof given by Minakawa in order to obtain further informations on the conjugating homeomorphism and to compute the rotation numbers of f and g.

Definitions.

- A homeomorphism $f \in PL^+(S_r)$ has the **(D)-property** if the product of the f-jumps on each orbit is trivial. Or equivalently the break set $BP(f)$ is contained in the union of finitely many pieces of orbits $C_i = \{a_i, f(a_i), ..., f^l(a_i)\}$, $i \in I$ such that $\prod_{c \in C_i} \sigma_f(c) = 1$.

- For f with (D)-property, we define $\pi(f) = \prod_{i \in I,\ 0<k\leq l_i} \sigma_{f^{N+1}}(f^k(a_i))$, where $N = max\ l_i$ with l_i defined as above.

Lemma 1. *If $f, g \in PL^+(S_r)$ generate a freely acting group, then they have the (D)-property.*

Proof of Lemma 1. Let a be a break of f, we have to prove that the product of the f-jumps on the f-orbit of a is trivial. In fact it suffices to prove the triviality of the product of jumps on finite pieces of orbit $\{f^{-M}(a), ..., f^M(a)\}$, for any sufficiently large M.

Since f has finitely many break points and since the f-orbits of the points $g^k(a)$ are disjoints, it's possible to find $k \in \mathbb{N}$ such that the f-orbit of $g^k(a)$ doesn't meet $BP(f)$. Thus, for all $n \in \mathbb{Z}$, we have $\sigma_f(f^n(g^k(a)) = 1$. Futhermore, for all $x \in S_r$, we have :

$$\sigma_{g^k \circ f \circ g^{-k}}(g^k(x)) = \frac{\sigma_{g^k}(f(x))\sigma_f(x)}{\sigma_{g^k}(x)}.$$

Using the commutativity hypothesis, we have

$$1 = \sigma_f(f^n(g^k(a)) = \sigma_{g^k \circ f \circ g^{-k}}(f^n(g^k(a)) = \sigma_{g^k \circ f \circ g^{-k}}(g^k(f^n(a)) = \frac{\sigma_{g^k}(f(f^n(a))\sigma_f(f^n(a))}{\sigma_{g^k}(f^n(a))}.$$

It follows that $\quad \sigma_f(f^n(a)) = \dfrac{\sigma_{g^k}(f^n(a))}{\sigma_{g^k}(f^{n+1}(a))}, \quad$ hence the product

$$\prod_{n \in \mathbb{Z}} \sigma_f(f^n(a)) = \prod_{n \in \{-M,..,M\}} \sigma_f(f^n(a)) \qquad \text{telescopes to} \quad \frac{\sigma_{g^k}(f^{-M}a))}{\sigma_{g^k}(f^{M+1}(a))}.$$

Finally, by choosing M sufficiently large (depending on k) such that $f^{-M}(a)$ and $f^{M+1}(a)$ are not breaks of g^k, we get that the product of the f-jumps along the f-orbit of a is trivial. We conclude that f (and also g) has the (D)-property.

Lemma 2. *Let $f \in T_{r,\Lambda,A}$ with (D)-property and irrational rotation number. Then there exists a PL map H_f that conjugates f to the S_r Boshernitzan example $\mathcal{H}_r \circ B^{\Pi(f)}_{\rho(f)} \circ \mathcal{H}_r^{-1}$. Furthermore :*

If $\Pi(f) \neq 1$, then $\rho(f) = \frac{\log \alpha}{\log \Pi(f)}$, where $\alpha \in \Lambda$.

If $\Pi(f) = 1$, $\rho(f) = \frac{\alpha}{\beta}$, where $\alpha, \beta \in \Lambda$.

The break set $BP(H) \subset A$, the jump sets $\sigma(H) \subset \Lambda$ and $\sigma(B^{\Pi(f)}_{\rho(f)}) \subset \Lambda$. But, in general, the slope sets $\Lambda(H)$ and $\Lambda(B^{\Pi(f)}_{\rho(f)})$ are not contained in Λ.

Proof of Lemma 2. By definition of the (D)-property, the set $\ BP(f) \subset \bigsqcup_{i \in I} \mathcal{C}_i\ $ with $\mathcal{C}_i = \{f^k(a_i),\ k = 0, ..., l_i\}$ and $\prod_{c \in \mathcal{C}_i} \sigma_f(c) = 1$.

Consider a PL homeomorphism $H_f = H$ of S_r such that $f(a_i)$, ..., $f^{l_i}(a_i)$ for $i \in I$ are breaks of H with associated jumps $\sigma_H(f^k(a_i)) = \sigma_{f^{N+1}}(f^k(a_i))$ for $k = 1, ...l_i$, where $N = max\ l_i$. Note that for a_i we also have $\sigma_H(f^k(a_i)) = \sigma_{f^{N+1}}(f^k(a_i)) = 1$.

A necessary and sufficient condition for H to have these breaks and no more is that the product of their H-jumps is trivial, that is $\Pi(f) = 1$.

So, when $\Pi(f) = 1$, we normalise H by $H(0) = 0$.

So, when $\Pi(f) \neq 1$, we add a break point c such that $c \in A \setminus \cup\{a_i, ..., f^{l_i}(a_i)\}$ and $\sigma_H(c) = (\Pi(f))^{-1}$ and normalise H by $H(c) = 0$.

By definition, this map H has breaks in A, jumps in Λ. The slopes of H are in $frac(A) = \{\frac{a}{b} :\ a, b \in A\}$ because are ratio of lengths of two intervals with endpoints in $BP(H) \subset A$.

Now we have :

- $BP(H \circ f \circ H^{-1}) \subset \{H(a_i), ..., H(f^{l_i}(a_i)), i \in I\} \cup \{H(c), H(f^{-1}(c)\}$,
- for $i \in I$, $0 \leq k \leq l_i$, the jump :

$$\sigma_{H \circ f \circ H^{-1}}(H(f^k(a_i)) = \frac{\sigma_H(f^{k+1}(a_i)) \times \sigma_f(f^k(a_i))}{\sigma_H(f^k(a_i))} = \frac{\sigma_{f^{N+1}}(f^{k+1}(a_i)) \times \sigma_f(f^k(a_i))}{\sigma_{f^{N+1}}(f^k(a_i))} = 1,$$

- the jump $\sigma_{H \circ f \circ H^{-1}}(H(c)) = \Pi(f)$,
- the jump $\sigma_{H \circ f \circ H^{-1}}(H(f^{-1}(c)) = \Pi(f)^{-1}$.

Thus, when $\Pi(f) \neq 1$, the map $F = H \circ f \circ H^{-1}$ has exactly two breaks $0 = H(c)$ and $F^{-1}(0) = H(f^{-1}(c))$, it's a Boshernitzan on S_r ; its jump at the initial break 0 is $\Pi(f)$ and its rotation number is $\rho(f)$ but also $\dfrac{\log DF_+(0)}{\log DF_+(0) - \log DF_-(0)} = \dfrac{\log DF_+(0)}{\log \Pi(f)}$ as it's a Boshernitzan.

Moreover, $DF_+(0) = D_+(H \circ f \circ H^{-1})(H(c)) = \dfrac{DH(f(c))Df(c)}{DH_+(c)} = Df(c)\dfrac{DH(f(c))}{DH_+(c)} \in \Lambda$, as $Df(c) \in \Lambda$ and the ratio of the derivatives of H at two points is a product of H-jumps.

So $\rho(f) = \rho(F) = \dfrac{\log \alpha}{\log \Pi(f)}$, with α and $\Pi(f)$ in Λ.

Now, when $\Pi(f) = 1$, the map $F = H \circ f \circ H^{-1}$ is PL and has no break so it's a rotation of angle $\rho(f)$ but also of angle $F(0) = H \circ f \circ H^{-1}(0) = H(f(0)) = DH(f(0)).f(0) + \beta \in fracA$, as $DH(f(0) \in fracA$ and $\beta \in fracA$ (this can easily be etablished by writing that H is continuous at its breaks $\alpha_i \in A$, that $H(0) = 0$ and that the slopes are in $fracA$). This ends the proof of Lemma 2.

Remarks on Lemma 2.

- When Λ is a subset of $\mathbb{Q}$, the case $\Pi(f) = 1$ can't occur, since $\rho(f)$ would be rational.
- If f and g satisfy the hypothesis of lemma 2 and commute then $\pi(f) = \pi(g)$, since the conjugation to rotation with $h(0) = 0$ is $R_{H_f^{-1}(0)} \circ H_f \circ h_{\pi(f)} = R_{H_g^{-1}(0)} \circ H_g \circ h_{\pi(g)}$ and an easy calculation gives that $h_{\pi_1} \circ h_{\pi_2}^{-1}$ is PL if and only if $\pi_1 = \pi_2$.
- For similar reasons, if f and g satisfy the hypothesis of lemma 2 and are PL conjugate then $\pi(f) = \pi(g)$.

Proof of Theorem 2.B. By absurb, we suppose that there is $\mathbb{Z}^p \subset T_{r,(n_1,...,n_p)}$ acting freely.

If $p = 1$, this is impossible by theorem 1 A.

If $p \geq 2$, there exist p commuting homeomorphisms f_i of irrational rotation numbers $\rho(f_i)$ affinely $\mathbb{Q}$-independent. Using lemma 1 and 2 (and its remarks) we get $\rho(f_i) = \frac{\log \alpha_i}{\log \Pi(f_i)}$.

Furthermore all $\Pi(f_i)$ are egal to a same Π (the f_i commute) and the $\rho(f_i) = \frac{\log \alpha_i}{\log \Pi}$ and 1 must be $\mathbb{Q}$-independent. This means (multiplicating by $\log \Pi \in \mathbb{Q}$) that the $\log \alpha_i$ and $\log \Pi$ are $\mathbb{Q}$-independent. But, $\log \alpha_i$ and $\log \Pi$ are $(p + 1)$ real numbers in the additve group generated by the p real numbers $\log n_i$, so they are $\mathbb{Q}$-dependent, a contradiction.

Proof of Theorem 2.C. If $f \in T_{r,(n_i)}$ is contained in a $\mathbb{Z}^2$ acting freely, by lemma 1, the map f has (D)-property. Thus, by lemma 2 (and its remarks), the rotation number of f is $\frac{\log \alpha}{\log \Pi(f)}$ with α and $\Pi(f)$ in $< n_i >$. This proves the claim.

Proof of Theorem 2.C'. Fix $p \geq 1$ and fix $\rho = \frac{\log \alpha}{\log \beta} \in]0,1[$ with $\alpha,\ \beta$ in $\Lambda =< n_i >$ and $1 < \alpha < \beta$. As $\beta - 1 \in (1-\Lambda)A = (d\mathbb{Z})\Lambda$ we can write $\beta - 1 = dn_\beta\lambda$ with $n_\beta \in \mathbb{Z}$ and $\lambda \in \Lambda$.

Let $r = dn_\beta$ and consider the Boshernitzan f on S_r defined by its two slopes

$$\lambda_1 = \alpha \text{ and } \lambda_2 = \frac{\lambda_1}{\beta} = \alpha\beta^{-1} < 1.$$

For proving that f is in $T_{r,(n_i)}$, it suffices to prove that its break $a_i \in \mathbb{Z}[\frac{1}{m}]$. We have $a_i = r\dfrac{1-\lambda_1}{\lambda_2-\lambda_1} = dn_\beta \times \dfrac{\alpha - 1}{\alpha\beta^{-1}(\beta-1)} = \dfrac{\alpha-1}{\lambda\alpha\beta^{-1}} \in A$. The rotation number of f is $\rho(f) = \frac{\log \lambda_1}{\log \lambda_2 - \log \lambda_1} = \frac{\log \alpha}{\log \beta}$. But $r = dn_\beta$ equal d $(mod\ d)$ so using Bieri-Strebel criterion the group $T_{d,(n_i)}$ contains a PL-conjugate to f that is a map of rotation number $\frac{\log \alpha}{\log \beta}$.

Remark also that for $p = 1$, the result is the consequence 2 of theorem 1.

3.5 Proof of the consequences of theorem 2.

Consequence 1 is a reformulation of A and B. The consequences 2, 3 and 4 use easy technical lemmas (lemma 4 and 5) and a **Brin version ([4]) of the Rubin theorem of [21] :**

Let G be a group and ϕ, φ be two representations of G in $Homeo^+(S^1)$. If the subgroups $\phi(G)$ and $\varphi(G)$ are both locally dense then ϕ and φ are topologically conjugate (that is there exists $h \in Homeo(S^1)$ such that $\phi(g) = h \circ \varphi(g) \circ h^{-1}$, $\forall g \in G$).

Where a subgroup H of $Homeo^+(S^1)$ is said to be **locally dense** *if for all open subset U of S^1, the group $G_U = \{g \in H : supp\ g \subset U\}$ has every orbit in U locally dense (i.e dense in some non empty open subset of U).*

Lemma 3. *Fix two integers $r \geq 1$ and $k \geq 2$, then the group $T_{r,k}$ is a locally dense subgroup of $Homeo^+(S^1)$.*

Proof of lemma 3. Let U be an open subset of S^1 and $x_0 \in U$. We have to prove that the closure $cl(G_U(x_0))$ of $G_U(x_0)$ contains an open subset of U. Let's choose $a_0, b_0 \in A = \mathbb{Z}[\frac{1}{k}]$ such that $x_0 \in I_0 =]a_0, b_0[\subset U$. We'll prove that $cl(G_{I_0}(x_0))$ contains a non empty open subset.

For $\alpha \in]0, (k-1)min(x_0 - a_0, \frac{b_0 - x_0}{k})[\cap A$, we consider the PL-homeomorphism f_α defined by :

$f_\alpha(x) = kx + (1-k)a_0$ if $x \in [a_0, a_0']$ with $a_0' = a_0 + \frac{\alpha}{k-1}$,

$f_\alpha(x) = x + \alpha$ if $x \in [a_0', b_0']$ with $b_0' = b_0 - \frac{k\alpha}{k-1}$,

$f_\alpha(x) = \dfrac{1}{k}x + (\dfrac{k-1}{k})b_0$ if $x \in [b_0', b_0]$,

$f_\alpha(x) = x$ elsewhere.



Fig. 2

We have $f_\alpha \in T_{r,k}$, its support $suppf_\alpha = [a_0', b_0'] \subset I_0 \subset U$ and $f_\alpha(x_0) = x_0 + \alpha$ for all α in the dense set $A' =]0, (k-1)min(x_0 - a_0, \frac{b_0 - x_0}{k})[\ \cap A$ as $a_0 \leq a_0' < x_0 < b_0' \leq b_0$.

Thus, cl($\{f_\alpha(x_0), \alpha \in A'\}$) has non empty interior and is contained in the interior of $cl(G_{I_0}(x_0))$. This ends the proof of lemma 3.

Consequences : The groups $T_{r,(n_i)}$ are locally dense subgroups (since they contain some $T_{r,k}$). **Each isomorphism between such groups and each automorphism of such groups is realized by a conjugation in Homeo (S^1).**

Lemma 4 : *Let* $\Gamma_1 =< f_1, g_1 > \subset Homeo^{\,+}(S_{r_1})$ *and* $\Gamma_2 =< f_2, g_2 > \subset Homeo^{\,+}(S_{r_2})$ *be two* $\mathbb{Z}^2$ *that act freely on circles and such that* $\rho(f_1) = \rho(f_2)$ *and* $\rho(g_1) = \rho(g_2)$*. Then* Γ_1 *and* Γ_2 *are conjugate, and the maps such that* $\Gamma_2 = h \circ \Gamma_1 \circ h^{-1}$ *are*

$$h_b = H_{f_2}^{-1} \circ \mathcal{H}_{r_2} \circ h_{\pi(f_2)} \circ R_b \circ h_{\pi(f_1)}^{-1} \circ {\mathcal{H}_{r_1}}^{-1} \circ H_{f_1} \quad \textit{with } b \in S^1,$$

where H_f *is the map defined in lemma 2,* $\mathcal{H}_r$ *is the homothety of ratio* r *and* h_π *is the conjugation between the Boshernitzan* B_ρ^π *and the rotation by* ρ*.*

The proof of lemma 4 is the combination of lemma 2 and properties of Boshernitzan-examples and we leave it to the reader.

Lemma 5 : *Let* $\pi_1,\ \pi_2$ *in* $]0,1]$ *and* a, b *in* S^1*, denote* $F_b = h_{\pi_2} \circ R_b \circ h_{\pi_1}^{-1}$*, then* $F_b \circ R_a \circ F_b^{-1}$ *is a PL-map if and only if* $a = 0$ *or* $\pi_1 = 1$ *or* $\pi_1 = \pi_2$*.*

Proof of lemma 5. The calculus are done in $[0,1[$, circle homeomorphisms f being identified with bijections $\tilde{f}$ (mod 1) of $[0,1[$.

We write $h_\pi(x) = \frac{\pi^x - 1}{\pi - 1} = \frac{1}{\pi-1}(e^{x \log \pi} - 1)$ et $h_\pi^{-1}(x) = \frac{1}{\log \pi} \log(x(\pi - 1) + 1)$.

We compute $F_b(x) = h_{\pi_1} \circ R_b \circ h_{\pi_2}^{-1}(x) = \dfrac{1}{\pi_1 - 1} \left(\pi_1^{b(x)} (x(\pi_2 - 1) + 1)^{\frac{p}{s}} - 1 \right)$, where $p = \log \pi_1,\ \ s = \log \pi_2$ and $b(x) \in \{b, b-1\}$.

Finally, $F_b \circ R_a \circ F_b^{-1}(x) = \dfrac{1}{\pi_2 - 1} \left(\pi_2^{-b_1(x)} \left(a(\pi_1 - 1) + \pi_1^{b_2(x)} (x(\pi_2 - 1) + 1)^{\frac{p}{s}} \right)^{\frac{s}{p}} - 1 \right)$, where $b_1(x), b_2(x) \in \{b, b-1\}$.

Finally, $F_b \circ R_a \circ F_b^{-1}(x)$ is PL if and only if $a'(\pi_1 - 1) = 0$ or $\frac{p}{s} = 1$, this gives lemma 5.

Proof of consequence 2 : “ Topological rigidity implies PL rigidity”.

Let ϕ be a representation of $G = T_{r,(n_i)}$, $p \geq 3$ in $PL^+(S_1)$ that is topologically conjugate by h to the standard one. By eventually composing h with the standard involution $(x \mapsto 1 - x)$, we may assume that h is orientation preserving. Consider a rank 2 subgroup $< f_1, g_1 >$ contained in the “free” $\mathbb{Z}^{p-1}$ subgroup of G that has been constructed in the previous section (proof of theorem 2.A). Changing r in its class mod d, we may assume that f_1 is a Boshernizan : $f_1 = \mathcal{H}_{r_1} \circ h_{\pi_1} \circ R_{\rho_1} \circ h_{\pi_1} \circ \mathcal{H}_{r_1}$.

The image of $< f_1, g_1 >$ is a $\mathbb{Z}^2$ acting freely, we can apply lemma 4 for $< f_1, g_1 > \subset G$ and $< \phi(f_1), \phi(g_1) > \subset \phi(G)$. We obtain that h is one of the h_b. Let $R_a \in G$ be a non trivial rotation, we have $\phi(R_a) = h_b \circ R_a \circ h_b^{-1} \in PL^+(S^1)$. Replacing h_b by its expression, it means that $H_{f_2}^{-1} \circ h_{\pi(f_2)} \circ R_b \circ h_{\pi(f_1)}^{-1} \circ \mathcal{H}_{r_1}^{-1} \circ R_a \circ \mathcal{H}_{r_1} \circ h_{\pi(f_1)} \circ R_b^{-1} \circ h_{\pi(f_2)}^{-1} \circ\ H_{f_2} \in PL^+(S^1)$.

Using the fact that H_f are PL-maps and lemma 5, we get that $\pi(f_1) = 1$ that is impossible as f_i is a rational homeomorphism with irrational rotation number or $\pi(f_1) = \pi(f_2)$ in this case $h_{\pi(f_2)} \circ R_b \circ h_{\pi(f_1)}^{-1}$ is a Boshernitzan so it's PL. Finally, we have proved that h_b is PL.

Proof of consequence 3 : “ Non exoticity of automorphisms and isomorphisms”.

According to Rubin theorem, isomorphisms and automorphisms of Thompson-Stein groups are realized by topological conjugations. According to the previous point, they are realized by PL-conjugation, provided the fact that $p \geq 3$. More precisely, it's possible to give a complete **description of PL-automorphisms.** This can be found in [1] or in [16] and prove that a PL automorphism of $G = T_{r,\Lambda,A}$ has slopes in $invA$ and breaks in A.

Proof of consequence 4 : “Distincts Thompson-Stein groups are not isomorphic”.

Suppose, by absurd, that $G = T_{r,(n_i)}$ and $G' = T_{r,(n'_j)}$ are isomorphic and G is of free rank at least 2. By consequence 3, the groups G and G' are PL-conjugate. Using the summary at the end of proof of theorem 2A, for all $i \in \{1, .., p\}$ the group $G = T_{r,(n_i)}$ contains a map f_i with rotation number $\rho(f_i) = \frac{\log n_i}{\log \Pi}$. The image f'_i of f_i in $G' = T_{r,(n'_i)}$ is contained in a free

$\mathbb{Z}^2$ so its rotation number $\rho(f'_i) = \frac{\log \alpha'}{\log \Pi(f'_i)}$, $\alpha' \in < n'_j >$ and as f'_i is PL conjugate to f_i we also have $\Pi(f'_i) = \Pi$ (according to remarks on lemma 2). So, for all $i \in \{1, ..., p\}$ the integer $n_i \in < n'_j >$ and conversely. Consequently, the groups $< n_i > = < n'_j >$ and eventually changing the basis of the slope sets we have $n_i = n'_j$, hence the groups $G = G'$.

4 Proof of theorem 3.

4.1 Dynamical properties for representations of $\mathbf{G = T_{1,(n_i)}}$.

In this section, we study the dynamics of a non trivial representation $\phi : G \to Homeo^+(S^1)$.

Notations. $F = \{g \in G : g(0) = 0\}$,
$F' = [F, F] = \{g \in G : g(0) = 0 \text{ and } Dg(0) = 1\}$,
$A = \mathbb{Z}[\frac{1}{m}]$, where $m = lcm(n_i)$ is identified to the subgroup of G consisting of rotations i.e the group of rotations of angles in A.

Algebraic properties, useful for dynamical properties on F and G :

- F doesn't contain any free group of rank 2 (Brin-Squier) and has no torsion.
- F' is simple (Bieri-Strebel, Stein).
- $T_{1,k}$, for all $k \geq 2$ integer in $< n_i >$, is simple (Brown), is contained in G and contains A.

Proposition 1. The group $\phi(F)$ acts on S^1 with a finite orbit.

Proof of proposition 1. By a Margulis theorem [Ma 2000], since according to Brin-Squier F contains no free group of rank 2, the group $\phi(F)$ preserves a measure μ on S^1. Thus, the rotation number map ρ is a representation of F into an abelian group. In particular, it's trivial on the first derivated group $\phi([F, F])$ and consequently, any element of $\phi([F, F])$ has a fixed point.

The **dynamical alternatives theorem** *(see [Gh2000] prop. 5.6) says that $\phi(F)$ satisfies one of the mutually exclusive following possibilities :*
1. there is a finite orbit or
there is a unique minimal K that is the accumulation set of any orbit and
2. $K = S^1$ and all orbits are dense or
3. $K \neq S^1$ is said to be exceptionnal (K is homeomorphic to the Cantor set).

If $\phi(F)$ has no finite orbit, the support $supp\mu$ of μ coincide with the unique minimal K of $\phi([F, F])$. If $K = S^1$ then by setting $h(x) = \mu([0, x])$, one constructs a conjugation from $\phi(F)$ to a group of rotations and F is an abelian group, this is not the case.

So, $K = supp\mu$ is a cantor set and any point in K is fixed by $\phi([F, F])$.

Moreover, let g_i, $i = 1, ..; p$ be the generators of F, they have rational rotation numbers $\frac{p_i}{q_i}$, since if not by Denjoy et unique ergodicity μ would be of total support.

Then the finite type abelian group $H = \frac{F}{[F,F]}$ acts on K (since $[F, F] \subset Ker\phi_{|K}$) and any generator g_i is of finite order q_i. So for all $x \in K$, we have $H(x) = \{g_1^{s_1} .....g_p^{s_p}(x), 0 \leq s_i < q_i\}$ that is finite.

Finally, the $\phi(F)$ orbit of any $x \in K$ is the set of the $\phi(f).x_0 = \gamma c(x_0)$, where $c \in \phi([F, F])$, $\gamma \in \{g_1^{s_1} .....g_p^{s_p}, 0 \leq s_i < q_i\}$, so it's finite since $\gamma c(x_0) = \gamma(x_0)$. This contadicts the dynamical alternatives theorem. This ends the proof of the proposition 1.

Proposition 2. *The groups $\phi(G)$, $\phi(A)$ and $\phi(T_{1,k})$ for k integer in $< n_i >$ have no finite orbits and have a unique same minimal K that is the accumulation set of any orbit.*

Proof of Proposition 2.

Step 1 : the groups $\phi(G)$, $\phi(A)$, $\phi(T_{1,k})$ for k integer in $< n_i >$ have no finite orbits.

The group $\phi(\mathbb{Z}[1/k])$ is isomorphic to $\mathbb{Z}[1/k]$ by proposition 2, so it contains element of arbitrary big order, so it has no finite orbits. Now, the groups $\phi(G)$ and $\phi(T_{1,k})$ contain some $\phi(\mathbb{Z}[1/k])$, so they also have no finite orbits.

Step 2 : the groups $\phi(G)$, $\phi(A)$, $\phi(T_{1,k})$, for k integer in $< n_i >$ have the same unique minimal K.

According to dynamical alternatives theorem, each of these groups $\phi(H)$ have a unique minimal denote by $K(H)$. From inclusions between these groups, follow the inclusions :

$K(\mathbb{Z}[1/k]) \subset K(T_{1,k}) \subset K(G)$, k integer in $< n_i >$
$K(\mathbb{Z}[1/k]) \subset K(A) \subset K(G)$.

We first prove that $K(A) = K(\mathbb{Z}[1/k])$. –For any $a \in A$ and any $\kappa \in \mathbb{Z}[1/k]$, we have $R_\kappa \circ R_a(K(\mathbb{Z}[1/k])) = R_a \circ R_\kappa(K(\mathbb{Z}[1/k])) = R_a(K(\mathbb{Z}[1/k]))$. This means that the set $R_a(K(\mathbb{Z}[1/k]))$ is $\mathbb{Z}[1/k]$-invariant, so by minimality and unicity of $K(\mathbb{Z}[1/k])$ this set contains $K(\mathbb{Z}[1/k])$.

As the same holds for $(-a) \in A$, we get $K(\mathbb{Z}[1/k])$ is A-invariant, so $K(\mathbb{Z}[1/k])$ contains $K(A)$ by minimality and unicity, this is the missing inclusion.–

Now, we prove that $K(A) = K(G)$. –As each $g \in G$ can be written as $g = R_a \circ f$, with $f \in F$, $a \in A$; we have $\phi(g) = \phi(a) \circ \phi(f)$. But we have proved that there exists a point x_0 with a finite $\phi(F)$-orbit : $\{x_0, ..., x_p\}$, so $\phi(g).x_0 \in \bigcup_{i=0}^{p} \phi(a).x_i$, for all $g \in G$. Thus $\phi(G).x_0 = \bigcup_{i=0}^{p} \phi(A).x_0$ and so on for the accumulation sets that are exactly the $K(\ )$.

Hence, we have $K(G) = K(A)$.–

Finally, we have proved that $K(\mathbb{Z}[1/k]) = K(A) = K(G)$ and using the first inclusions we have that this set is also $K(T_{1,k})$. According to the dynamical alternatives theorem, this set is the accumulation set of any orbit.

4.2 Proof of theorem 3.A, from now $\mathbf{G = T_{1,(2,n_2,\dots,n_p)}}$, $\mathbf{p \geq 2}$.

Step 1. Any non trivial representation ϕ of G in $\mathrm{PL}^+(S^1)$ or $\mathrm{Diff}^2_+(S^1)$ such that $\phi(G)$ has dense orbits is conjugate to the standard PL representation.

Proof – The induced action on the classical Thompson group $T_{1,2} \subset G$ satisfies the hypothesis of III proposition 3.9 p 230 in [11] : it's non trivial and the group $\phi(\mathbb{Z}[1/2])$ has dense orbits because it has the same minimal than $\phi(G)$. Thus, by Ghys-Sergiescu results, $\phi(T_{1,2})$ is conjugate to the PL standard action $T_{1,2}$. So both actions G and $\phi(G)$ satisfy the hypothesis of Rubin-Brin result, they are topologically conjugate.–

Step 2. Any non trivial representation ϕ of G in $\mathrm{Homeo}^+(S^1)$ is semi-conjugate to a representation ϕ_1 of G in $\mathrm{Homeo}^+(S^1)$ such that $\phi_1(G)$ has dense orbits.

Proof – If $K(G)$ is S^1, the representation $\phi(G)$ has dense orbits. If $K(G)$ is not S^1, it's a Cantor set, the devil stairs associated to K semi-conjugates ϕ to ϕ_1 with dense orbits.–

It's noted in [11]–proof of III Theorem 3.12 p 233– that when ϕ is C^2 or PL, the action ϕ_1 failed not to be so regular but it satisfies all thinks that are needed for proving that $\phi_1(T_{1,2})$ is conjugate to the PL standard action $T_{1,2}$ which implies in our case that both actions G and $\phi_1(G)$ satisfy the hypothesis of Rubin-Brin result, and hence are topologically conjugate. We are now able to give

the end of the proof of theorem 3.A. Let ϕ be a representation of G in $PL(S^1)$ or Diff (S^1), composing ϕ with the standard involution $(x \mapsto 1-x)$ we can suppose that ϕ is orientation preserving. If $\phi(G)$ has dense orbits then it is conjugate to the standard PL representation, by step 1. If not, by step 2 and the previous remark, ϕ is semi-conjugate to such an action ϕ_1, so it's semi-conjugate to the standard PL action. But, as $p \geq 2$, the group G contains at least one element of irrational rotation number for which Denjoy theorem applies, this and unique ergodicity implies that the semi-conjugation is a conjugation.

4.3 Proof of theorem 3.B.

Consider the Thompson-Stein group $G = T_{1,(2,n_2,..n_p)}$ with $p \geq 2$. By the theorem 2C', the group G contains an homeomorphism B of irrational rotation number $\alpha = \frac{\log 2}{\log n_2}$ a diophantine number (see [21]) of irrationality measure β (for example for $n_2 = 3$ we have $\beta = 7,616....$)

Suppose that there exists a representation $\phi : G \to \text{Diff}^{\,k}(S^1)$, for some $k \geq 2$. By theorem 3.A, $\phi(G)$ is topologically conjugate by some homeomorphism h to G.

So $\phi(B) = h \circ B \circ h^{-1}$ is a C^k-diffeomorphism with diophantine rotation number α of irrationality measure β, by Katznelson-Ornstein result ([15]) it's C^1-conjugate to R_α provided $k \geq \beta + 1$.

Thus for $k \geq \beta + 1$, there exists a C^1-diffeomorphism H such that $R_\alpha = H \circ h \circ B \circ h^{-1} \circ H^{-1}$. As B is PL-conjugate to a Boshernitzan of rotation number α, it is piecewise C^∞ conjugate to R_α. Using the unique ergodicty of R_α, we get that the map $H \circ h$ is piecewise C^∞, so we get that h is piecewise C^1 with a finite number N of breaks.

Finally, every $g \in G$ can be written as $g = h^{-1} \circ \phi(g) \circ h$ so it is piecewise C^1 and has at most $2BP(h) = 2N$ breaks, this is impossible as G contains elements with arbitrary large number of breaks.

4.4 Proof of the consequence of theorem 3.

This is an immediate consequence of theorem 3A and consequence 2 of theorem 2 ("topological rigidity implies PL rigidity").

5 References.

1. R. Bieri and R. Strebel, *On groups of PL homeomorphisms of the real line.* Preprint, Math. Sem. der Univ. Frankfurt, Frankfurt am Main (1985).
2. M. Boshernitzan, *Dense orbits of rationals.* Proc. AMS, **117** (4), 1201-1203 (1993).
3. M. G. Brin, *The chameleon groups of Richard J. Thompson : Automorphisms and dynamics*, Inst. Hautes Études Sci. Publ. Math. **84**, 5–33 (1996).
4. M. G. Brin, *Higher dimensionnal Thompson's groups.* Geom. Dedicata **108**, 163–192 (2004).
5. M. G. Brin, C. Squier, *Groups of piecewise linear homeomorphisms of the real line.* Invent. Math. **79** (3), 485–498 (1985).
6. M. G. Brin, F. Guzmán, *Automorphisms of generalized Thompson groups.* J. Algebra **203**(1), 285–348 (1998).
7. K. S. Brown, *Finiteness properties of groups.* J. Pure Appl. Algebra **44**, 45–75. (1987).
8. J. Cannon, W. Floyd, W. Parry, *Introductory notes on Richard Thompson's groups.* L'Ens. Math. **42**, 215-256 (1996).
9. A. Denjoy, *Sur les courbes définies par les équations différentielles à la surface du tore.* J. Math. Pures Appl. **11**, 333-375 (1932).
10. D.S Farley, *Proper isometric actions of Thompson's groups on Hilbert space.* Int. Math. Res. Not. **45**, 2409–2414 (2003).
11. É. Ghys, V. Sergiescu, *Sur un groupe remarquable de difféomorphismes du cercle.* Comm. Math. Helv. **62**, 185-239 (1987).
12. É. Ghys, *Groups acting on the circle.* L'Ens. Math. **62**, 215-256 (2000).
13. M. Herman, *Sur la conjugaison différentiable des difféomorphismes du cercle à des rotations.* Inst. Hautes Etudes Sci. Publ. Math. **49**, 5-234 (1979).
14. G. Higman, *Finitely presented infinite simple groups.* Notes on Pure Mathematics **8** (1974). Department of Pure Mathematics,Department of Mathematics, I.A.S. Australian National University, Canberra, 1974. vii+82 pp
15. Y. Katznelson, D. Ornstein, *The differentiability of the conjugation of certain diffeomorphisms of the circle.* Erg. Th. and Dyn. Syst. **9**, 643-680 (1989).
16. I. Liousse, *Nombre de rotation dans les groupes de Thompson généralisés, applications.* http ://hal.ccsd.cnrs.fr/ccsd-00004554.
17. G. Margulis, *Free subgroups of the homeomorphism group of the circle.* C. R. Acad. Sci. Paris Sér. I Math. **331**(9), 669–674 (2000).
18. H. Minakawa, *Classification of exotic circles of* $\mathrm{PL}_+(S^1)$. Hokkaido Math. J. **26** (3), 685–697 (1997).
19. H. Poincaré, *Oeuvres complètes.* **t.1**, 137-158.
20. G. Rhin, *Approximants de Padé et mesures effectives d'irrationnalité*, séminiare de théorie des nombres Paris 1985-1986, Prog. math **71**, 155-164 (1987).
21. Y. Rubin, *Locally moving groups and reconstruction problems*, Math. Appl. **354**, Kluwer Acad. Publ., Dordrecht (1996).
22. M. Stein, *Groups of piecewise linear homeomorphisms*, Trans. A.M.S. **332**, 477-514 (1992).

Laboratoire Paul PAINLEVÉ, U.M.R 8524 au CNRS
U.F.R. de Mathématiques, Université Lille I
59655 Villeneuve d'Ascq Cédex, France